


Contents list available at JMCS

Journal of Mathematics and Computer Science

Journal Homepage: www.tjmcs.com



Cyclic Edge Extensions – Self-centered Graphs

Medha Itagi Huilgol

Department of Mathematics, Bangalore
University, Central College Campus,
Bangalore, Karnataka, India

medha@bub.ernet.in

Chitra Ramaprakash

Department of Mathematics Bangalore
University, Central College Campus,
Bangalore, Karnataka, India

r.chitraprakash@gmail.com

Article history:

Received March 2014

Accepted April 2014

Available online April 2014

Abstract

The eccentricity $e(u)$ of a vertex u is the maximum distance of u to any other vertex of G . The maximum and the minimum eccentricity among the vertices of a graph G are known as the diameter and the radius of G respectively. If they are equal then the graph is said to be a self - centered graph. Edge addition /extension to a graph either retains or changes the parameter of a graph, under consideration. In this paper mainly, we consider edge extension for cycles, with respect to the self-centeredness(of cycles),that is, after an edge set is added to a self centered graph the resultant graph is also a self-centered graph. Also, we have other structural results for graphs with edge -extensions.

Keywords: Self centered graphs, Edge extension graphs, reduced radius, reduced diameter of cycles, Iterations of cycles and paths.

1. INTRODUCTION.

Unless mentioned otherwise, for terminology and notation the reader may refer Buckley and Harary [7], new ones will be introduced as and when found necessary.

In this paper, by a graph G , we mean a simple, undirected, connected graph without self-loops. The order and size are respectively the number of vertices denoted by p and the number of edges denoted by q .

The distance $d(u, v)$ between any two vertices u and v , of G , is the length of the shortest path between u and v . The eccentricity $e(u)$ of a vertex u is the distance to a farthest vertex from u . The maximum and the

minimum eccentricity amongst the vertices of G are respectively called the *diameter*, $diam(G)$ and *radius*, $rad(G)$. If $diam(G) = rad(G)$, then the graph G is said to be a *self-centered graph*. If $dist(u, v) = e(u)$, ($v \neq u$) then we say that v is an *eccentric* vertex of u .

Harary [8], introduced the concept of changing and unchanging of a graphical invariant i , with interest in determining those for which $i(G - v) = i(G)$ and $i(G - v) \neq i(G)$ for all vertices v of G , $i(G - e) = i(G)$ and $i(G - e) \neq i(G)$ for all edges e of G and $i(G + e) = i(G)$ and $i(G + e) \neq i(G)$ for all edges e of \bar{G} , the compliment of G . These concepts have been studied quite well for several invariants by Dutton et al. [9], Brigham et al. [15], [16], Harary [17], Lawson et al. [18], Medidi [19], Walikar et al. [11], [12], [13], [14] and Akram [4], [5]. Usually, these kind of studies reflect the variation of a parameter due to vertex or edge removal or edge addition, which find their applications in network analysis as they measure the results of link or equipment failure or network enhancement.

Janakiraman et al. [10] developed algorithms for constructing self-centered graphs from trees and connected graphs by adding edges. The authors defined a new concept called *center number* of a graph, denoted by $sc_r(G)$, which gives the minimum number of edges that can be added to a graph G to obtain a self-centered graph.

Akram [4] studied the addition of edges by introducing the concepts of *edge extension* set of graphs, *edge extensible class* of graphs and the *edge extensibility number* of a graph. He defined a non-empty set S of edges as *edge-extension set*, such that every edge in S joins two non-adjacent vertices in G . And

$G + S$, the graph after adding S to it is called the edge extension graph.

Definition 1 [4]

Let τ be a class of graphs satisfying certain property. Then τ is called edge extensible class, if for every graph $G \in \tau$, G is complete, or there exists an extension edge e such that $G + e \in \tau$.

Definition 2 [4]

Let G be a non-trivial simple graph (not complete). The simple graph obtained from G by adding a non-empty set of edges S such that every edge in S joins two non-adjacent vertices in G is called edge extension graph, and is denoted by $G + S$, S is called the *edge extension set*. In particular, if S consists of a single element e , then e is called the *extension edge*, and the graph is denoted by $G + e$.

We can see that the graph $G + S$ has the vertex set and the edge set as follows.
 $V(G + S) = V(G)$ and $E(G + S) = E(G) \cup S$.

Definition 3 [4]

Let τ be a class of graphs with certain property and $G \in \tau$ be non-trivial. The *edge extensibility number* of G with respect to τ is the smallest positive integer m , if exists, such that there exists an *edge extension set* S of cardinality ' m ' in such a way, the graph $G + S \in \tau$. We write $m = ext_\tau(G)$. If such a number does not exist for G , then we say that the corresponding edge extensibility number is ∞ .

It is easy to note that the class of connected graphs is edge extensible class, but not regular graphs. On the other hand a tree with respect to the class of trees τ has extensibility number ∞ . Akram [4], [5] has proved various results on extensibility of graphs and digraphs. Also, we can find results related to extension number of class of graphs viz., regular graphs, Eulerian graphs.

In this paper we consider self centered graphs as the collection τ and obtain edge extensibility number of some self centered graphs. Since cycles are the minimum sized self centered graphs, we find out

$ext_{\tau}(G) = m$.

It is clear that $m \neq 1$ for $\tau = C_p$. Hence, it is interesting and challenging to find the $ext_{\tau}(C_p)$.

2. Edge extension for cycles

In this section we consider edge extensions for cycles. As stated above it is clear that $m \neq 1$, with $m = ext_{\tau}(C_p)$ where τ is the class of self-centered graphs. When an edge (set) is added to a cycle we see that it does not remain a self-centered graph. So the first result discusses the minimum number of edges required to be added to a cycle such that the resulting graph is a self centered graph. And, we denote $m_i = ext_{\tau}(C_p)$ where i denotes the amount by which the radius(diameter) of the cycle C_p is reduced.

Lemma 2.1

Let C_p be the class of cycles of length (order) p . Let τ be the set of self-centered graphs. Then,

$$m_i = ext_{\tau}(C_p) = \begin{cases} 3, & \text{if } p \text{ is even and } p \geq 12, \\ 4, & \text{if } p \text{ is odd and } p \geq 11, \\ 2, & \text{when } p = 4, 8, 9, 10, \\ 5, & \text{when } p = 5, \\ 3, & \text{when } p = 6, 7. \end{cases}$$

Proof: Label the vertices of the cycle as $u_1, u_2, u_3 \dots u_p$.

For $p \geq 12$, with p even, join $u_1 u_3, u_1 u_{p-1}$ and $u_2 u_4$ so that the resultant graph is a self-centered graph of radius $(p/2) - 1$

For $p \geq 11$, with p odd, join $u_1 u_3, u_1 u_{p-1}, u_2 u_4$ and the fourth edge from $u_{p+1/2} u_{p+3/2}$, to get a self-centered graph with reduced diameter, by one.

For C_4 and C_5 it is clear that $ext_{\tau}(C_4) = 2$ and $ext_{\tau}(C_5) = 5$ respectively, as they result into K_4 and K_5 on addition of edges.

For C_6 , $ext_{\tau}(C_6) = 3$ and for C_7 , $ext_{\tau}(C_7) = 3$.

And for C_8, C_9, C_{10} , the extension number is 2 as we can add the edges $u_1 u_3$ and $u_1 u_{p-1}$.

Hence the proof. \square

Lemma 2.2

Let C_p denote the class of cycles of length (order) p , with $p \geq 11$. Then,

$$m_2 = ext_{\tau}(C_p) = \begin{cases} 4, & \text{if } p \text{ is even and } p \geq 10, \\ 5, & \text{if } p \text{ is odd and } p \geq 11, \\ 7, & \text{when } p = 7, \\ 4, & \text{when } p = 8, \\ 9, & \text{when } p = 9. \end{cases}$$

Proof: Label the vertices of the cycle as $u_1, u_2, u_3, \dots, u_p$.

For $p \geq 10$, with p even, join the edges $u_1 u_3, u_1 u_{p-1}, u_3 u_6$ and $u_5 u_8$, to get a graph G whose diameter is two less than that of C_p .

For $p \geq 11$, with p odd, join the edges $u_1 u_3, u_1 u_{p-1}, u_p u_{p-3}, u_3 u_6$ and $u_5 u_8$.

For $p = 7$, adding one more C_7 to the existing one reduces the diameter by 2. For $p = 8$, adding 4 edges

to each of their eccentric nodes reduces the diameter by 2. For $p = 9$, adding 9 more edges (one more C_9) reduces the diameter by 2. Hence the proof. \square

Lemma 2.3

Let C_p denote the class of cycles with $p \geq 14$. Then,

$$m_3 = \text{ext}_\tau(C_p) = \begin{cases} 6, & \text{if } p \text{ is even and } p \geq 14, \\ 8, & \text{if } p \text{ is odd and } p \geq 15, \end{cases}$$

Proof: Label the vertices of the cycle C_p as $u_1, u_2, u_3, \dots, u_p$.

If p is even, with $p \geq 14$, then join the edges $u_1 u_3, u_3 u_6, u_5 u_8, u_1 u_{p-1}, u_{p-1} u_{p-4}$ and $u_{p-3} u_{p-6}$, so that the resultant graph has its diameter reduced by 3.

If p is odd, with $p \geq 15$, then join $u_1 u_3, u_2 u_4, u_3 u_6, u_5 u_8, u_1 u_{p-1}, u_p u_{p-3}, u_{p-2} u_{p-5}$ and $u_{p-4} u_{p-7}$ to obtain the required result. Hence the proof. \square

Similarly we can add edges to the cycle to reduce the radius/diameter until we get a complete graph. The above results give reduction of eccentricity of each vertex to be reduced by one, two or three. As generalization seems difficult, to find extension number for cycles, to be in class of self-centered graphs, the above results help us to measure the number of edges to be added.

In the next results we do not add single edge, instead, we add paths, but the resultant graph is a self-centered graph. This way of approach is motivated by Buckley [6] in which he considered graphs under edge operation. Akira Saito et.al [1], [2], and [3] had considered properties of cycles of particular length. We combine both these approaches in the coming results.

Definition 4[6]

If $a \geq 4$, then $C_a *s P_b$ consists of the graph formed from C_a by joining two vertices u and v of C_a at distance b from one another by s additional paths of length b ($b > 1$).

Definition 5[6]

If $a \geq 4$ and $1 < d < b$, then $C_a *s P_b *P_d$ is the graph formed from $C_a *s P_b$ by joining the vertex u to a vertex w in C_a at distance d from u by an additional path of length d .

Lemma 2.4

Let C_p be a cycle of odd length, where $p \geq 7$. Then a path of length P_p concatenated with two eccentric vertices in a cycle results in a self-centered graph.

Proof: Case (1) Consider a cycle C_{7+4k} , which is of length - 7 modulo 4 where $k = 0, 1, 2, 3 \dots$

Let P_{2p} be a path, where $p \in \mathbb{Z}$ and $p \geq 2$.

On concatenating one end vertex of the path to any vertex say u of C_p and the other end vertex (of the path) to the eccentric vertex of u , say v , results in a self-centered graph. The length of the path varies depending on the radius of the cycle. Hence the path length $2p$ is one less or two less than the radius of the cycle.

Similarly, we can prove for the below two cases, by concatenating the specified path with a vertex and its eccentric vertex.

Case (2) Consider a cycle C_{9+4k} , which is of length $-9 \pmod{4}$ where $k = 0, 1, 2, 3 \dots$. Let P_{2p+1} be a path, where $p \in \mathbb{Z}$ and $p \geq 2$.

Case(3) Consider a cycle C_{4+2k} , which is of length $4 \pmod{2}$ where $k = 0, 1, 2, 3 \dots$. Let P_p be a path, where $m \in \mathbb{Z}$ and $m = \text{rad}(C_p - 1)$ or $m = \text{rad}(C_p - 2)$ or $m = \text{rad}(C_p)$.

Hence the proof. \square

Remark 1

A path of length P_2 can be added to C_7 and a path of length P_3 can be added to C_{11} to obtain a self centered graph. A path of length P_2 and P_3 can be added to C_9 and a path of length P_4 can be added to C_{13} to obtain a self centered graph.

In the next part we find the number of iterations, required for a cycle and a path to become a complete graph, using the concept of powers and the following three algorithms give the same.

Algorithm 2.1

In this algorithm we find the number of iterations required for cycle to be a complete graph. Let C_p be a cycle of length p .

STEP 1 : Input the cycle length p .

STEP 2 : Find the eccentricity of the given cycle by using $e = p/2$ if n is even or $e = \text{floor of } [p/2]$.

STEP 3 : If $e > 1$, then increase the iteration. Next we start adding edges for the next iteration such that the distance between any two vertices is less than or equal to the iterated power. This is done for all the vertices of the cycle.

STEP 4 : Again checking for the eccentricity of all the nodes. If $e_{\text{new}} = 1$, then GOTO STEP 6 else GOTO STEP 3.

STEP 5 : Print the iteration number.

STEP 6 : Print the number of iterations to get a complete graph.

STEP 7 : STOP.

Algorithm 2.2

In this algorithm we find the number of iterations required for path to be a complete graph. Let P_p be a path on $p + 1$ vertices.

STEP 1 : Input the path length.

STEP2 : Find the radius (minimum eccentricity) and the diameter (maximum eccentricity). The radius is denoted by $e_{\text{min}} = a$ and the diameter is denoted by $e_{\text{max}} = b$.

STEP 3 : If $e_{\text{min}} > 1$ and $e_{\text{max}} > 1$, increase the power else GOTO STEP 7

STEP 4 : Add edges to the consecutive node whose length \leq iterated power.

STEP 5 : Find the new eccentricities, $e_{\text{new-min}} = a/2$ and $e_{\text{new-max}} = b/2$

STEP 6 : If $a/2 = 1$ and $b/2 = 1$, GOTO STEP 7 else GOTO STEP 3.

STEP 7 : Print the number of iterations.

STEP 8 : STOP.

Algorithm 2.3

In this algorithm we find the iteration number for cycles to be self-centered by adding edges.
Let C_p be a cycle of length p .

STEP 1 : Input the cycle length p .

STEP 2 : Find the eccentricity of the given cycle by using $e = p/2$ if p is even or $e = \text{floor of } [p/2]$,
if p is odd.

STEP 3 : ' p ' denotes the length = p = edges .

STEP 4 : Input the eccentricity reduction value.

STEP 5 : From the first vertex start adding edges one by one such that the eccentricity of the other
vertices remain the same as inputted by the user.

STEP 6 : Perform STEP 5 until all the vertices have the same eccentricity.

STEP 7 : Check the eccentricity of the 2nd, 3rd and so on up to the p^{th} vertex. Check the eccentricity of
all the vertices. If they are same GOTO STEP- 8 or GOTO STEP 5.

STEP 8 : Output List of available vertices.

STEP 9 : Output list of edges added and the number of edges added.

STEP 10 : Output list of Invalid nodes where the edge addition is not possible.

STEP 11 : STOP.

CONCLUSION

Characterization on the number of edges to be added to a general graph seems to be difficult at this point of time. Hence, particular cases give insight about the edge additions to retain a particular property. The results discussed in this paper deal with additions done to a cycle to retain its self centeredness.

REFERENCES

Journal Articles

- [1] Akira Saito, Cycles of length 2 modulo 3 in graphs, Discrete Mathematics, 101(1992) 285-289.
- [2] Akira Saito, Gantao Chen, Graphs with a cycle of length divisible by three, Journal of Combinatorial theory,60(1994),No.2, 277-292.
- [3] Akira Saito, Linda Lesniak, Nathaniel Dean, Cycles of length 0 modulo 4 in graphs, Discrete Mathematics, 121(1993) 37-49.
- [4] Akram B. Attar, Edge Extension of Graphs and Digraphs, The Journal of Mathematics and Computer Science, 3(2011), No.1, 1-10.
- [5] Akram B. Attar, Extensibility of Graphs, Journal of Applied Mathematics, Islamic Azad University of Lahijan, 6(2009), No.21,1-9.
- [6] F. Buckley, Self-centered graphs with a given radius, Congressus Numerantium, 23(1979) 211- 215.
- [8] F. Harary, Changing and Unchanging radius of the graph, Linear Algebra and its Applications, 217(1995) 67-82.
- [9] Ronald D. Dutton, Sirisha R Medidi and Robert C Brigham, Changing and Unchanging of the Radius of a Graph, Linear Algebra and its Applications, 217(1995),67-82.
- [10] T.N. Janakiraman, M. Bhanumathi and S. Muthammai, Self - centered super graph of a graph and center number of a graph, Ars Combinatorica, 87(2008) 271-290.
- [11] H.B. Walikar, F. Buckley, M.K. Itagi, Radius - edge - invariant and diameter - edge- invariant graphs, Discrete Mathematics , 272(1)(2001) 119-126.

- [12] H.B. Walikar, F. Buckley, M.K. Itagi, Diameter essential edges in a graph, *Discrete Mathematics*, 259(2002) 211-225.
- [13] H.B. Walikar, F. Buckley, M.K. Itagi, Radius essential edges in a graph, *Journal of Combinatorial Mathematics and Combinatorial Computing*, 53(2005) 209-220.
- [14] H.B. Walikar, F. Buckley, Medha Itagi Huilgol, Diameter vital edges in a graph, *Aequationes Mathematicae*, 82(2011) 201-211.
- [15] R.C. Brigham and R.D. Dutton, An Extremal Problem for Edge Domination Insensitive Graphs, *Discrete Appl. Math*, 20(1988) 113-125.
- [16] R.C. Brigham and R.D. Dutton, Changing and Unchanging Invariants: The Edge Clique Cover Number, *Congr. Numer.*, 70(1990) 145-152.
- [17] F. Harary, Changing and Unchanging Invariants for Graphs, *Bull. Malaysian Math. Soc.*, 5(1982) 73-78.
- [18] L. M. Lawson and T.W. Haynes, Changing and Unchanging of the Node Covering Number of a Graph, *Congr. Numer.*, 77(1991) 157-162.

Books

- [7] F. Buckley, F. Harary, "Distance in Graphs", Addison Wesley, 1990.

Thesis Reference

- [19] S. Medidi, The Effect of Node Deletions, Edge Additions and Edge Deletions on Graphs: Node Connectivity, Edge Connectivity, Diameter, and Radius, Masters Project, Dept. of Computer Science, Univ. of Central Florida, 1992.